arXiv:1704.04925v1 [math.DG] 17 Apr 2017

VANISHING PROPERTIES OF p-HARMONIC ℓ-FORMS ON RIEMANNIAN MANIFOLDS

NGUYEN THAC DUNG AND PHAM TRONG TIEN

ABSTRACT. In this paper, we show several vanishing type theorems for p-harmonic ℓ-forms on Riemannian manifolds ($p \geq 2$). First of all, we consider complete non-compact immersed submanifolds M^n of N^{n+m} with flat normal bundle, we prove that any p-harmonic ℓ-forms on M is trivial if N has pure curvature tensor and M satisfies some geometric condition. Then, we obtain a vanishing theorem on Riemannian manifolds with weighted Poincaré inequality. Final, we investigate complete simply connected, locally conformally flat Riemannian manifolds M and point out that there is no nontrivial p-harmonic ℓ-form on M provided that Ric has suitable bound.

2000 Mathematics Subject Classification: 53C24, 53C21

Key words and phrases: p-harmonic functions; Flat normal bundle; Locally conformally flat; Weighted p-Laplacian; Weighted Poincaré inequality.

1.. INTRODUCTION

Suppose that M is a complete noncompact oriented Riemannian manifold of dimension n. At a point $x \in M$, let $\{\omega_1, \ldots, \omega_n\}$ be a positively oriented orthonormal coframe on $T_x^*(M)$, the Hodge star operator acting on the space of smooth ℓ-forms $\Lambda^\ell(M)$ is given by

$$*(\omega_{i_1} \wedge \ldots \wedge \omega_{i_\ell}) = \omega_{j_1} \wedge \ldots \wedge \omega_{j_{n-\ell}},$$

where $j_1, \ldots, j_{n-\ell}$ is selected such that $\{\omega_{i_1}, \ldots, \omega_{i_\ell}, \omega_{j_1}, \ldots, \omega_{j_{n-\ell}}\}$ gives a positive orientation. Let d is the exterior differential operator, so its dual operator δ is defined by

$$\delta = *d*.$$

Then the Hogde-Laplace-Beltrami operator Δ acting on the space of smooth ℓ-forms $\Omega^\ell(M)$ is of form

$$\Delta = -(\delta d + d\delta).$$

When M is compact, it is well-known that the space of harmonic ℓ-forms is isomorphic to its ℓ-th de Rham cohomology group. This is not true when M is non-compact but the theory of L^2 harmonic forms still have some interesting applications. For further results, we refer the reader to [Carron 1998, Carron 2007]. In [Li 1980], Li studied Sobolev inquality on spaces of harmonic ℓ-forms on compact Riemannian manifolds. He gave estimates of the bottom of ℓ-spectrum and proved that the space of harmonic ℓ-forms is of finite dimension provided that the Ricci curvature is bounded from below. In [Tanno 1996], Tanno studied L^2 harmonic ℓ-form on complete orientable stable minimal hypersurface M immersed in the Euclidean space $\mathbb{R}^{n+1}$. He showed that there are no non trivial L^2 harmonic ℓ-forms on M if $n \leq 4$. Later, Zhu generalized Tanno's results to manifolds with non-negative isotropic curvature (see [Zhu 2011]). He also proved in [Zhu 2001] that the Tanno's results are true if $M^n, n \leq 4$ be a complete noncompact strongly stable hypersurface with constant mean curvature in $\mathbb{R}^{n+1}$ or $\mathbb{S}^{n+1}$. Recently, in [Lin 2015a], Lin investigated spaces of L^2 harmonic ℓ-forms $H^\ell(L^2(M))$ on submanifolds in Euclidean space with flat normal bundle. Assumed that the submanifolds are of finite total curvature, Lin showed that the space $H^\ell(L^2(M))$ has finite dimension (See also [Zhu 2017] for the case $\ell = 2$). For further results in this direction, we refer

to [Lin 2015b, Lin 2015c, Lin 2016, Seo 2014, Zhu 2001, Zhu 2011] and the references therein. It is also worth to notice that the main tools to study the spaces of harmonic ℓ-forms are the Bochner formula and refined Kato type inequalities. In 2000, Calderbank et al. gave very general forms of Kato type inequalities in [Calderbank et al. 2000]. Then in [Wang 2002], Wang used them to prove a vanishing property of the space of L^2 harmonic ℓ-forms on convex cocompact hyperbolic manifolds. Later, in [Wan and Xin 2004], Wan and Xin studied L^2 harmonic ℓ-forms on conformally compact manifolds with a rather weak boundary regularity assumption. Recently, in [Cibotaru and Zhu 2012], Cibotaru and Zhu introduced a proof of the mentioned results from [Calderbank et al. 2000] avoiding as much as possible representation theoretic technicalities. The refined Kato type inequalities they obtained also refined the ones used in [Wang 2002, Wan and Xin 2004].

On the other hand, the p-Laplacian operator on a Riemannian manifold M is defined by

$$\Delta_p u := \mathrm{Div}(|\nabla u|^{p-2}\nabla u)$$

for any function $u \in W^{1,p}_{loc}(M)$ and $p > 1$, which arises as the Euler-Lagrange operator associated to the p-energy functional

$$E_p(u) := \int_M |\nabla u|^p.$$

Therefore, if u is a smooth p-harmonic function then du is a p-harmonic 1-forms. We refer the reader to [Moser 2007, Kotschwar and Ni 2009] for the connection between p-harmonic functions and the inverse mean curvature flow. Motivated by the above beautiful relationship between the space of harmonic ℓ-forms and the geometry of Riemannian manifolds, it is very natural for us to study the geometric structures of Riemannian manifolds by using the vanishing properties of p-harmonic ℓ-forms with finite L^q energy for some $p \geq 2$ and $q \geq 0$.

Recall that an ℓ-form ω on a Riemannian manifold M is said to be p-harmonic if ω satisfies $d\omega = 0$ and $\delta(|\omega|^{p-2}\omega) = 0$. When $p = 2$, a p-harmonic ℓ-form is exactly a harmonic ℓ-form. Some vanishing propeties of the space of p-harmonic ℓ-forms are given by X. Zhang in [Zhang 2001]. In particular, Zhang showed that there are no nontrivial p-harmonic 1-forms in $L^q(M), q > 0$ if the Ricci curvature on M is nonnegative. Motivated by Zhang's results, in [Chang et al. 2010], Chang-Guo-Sung proved that any bounded set of p-harmonic 1-forms in $L^q(M), 0 < q < \infty$, is relatively compact with respect to the uniform convergence topology. Recently, it is showed that the set of p-harmonic 1-forms has closed relationship with the connectedness at infinity of the manifold, in particularly, with p-nonparabolic ends. In [Dung and Seo 2017], the first author and Seo studied the connectedness at infinity of complete submanifolds by using the theory of p-harmonic function. For lower-dimensional cases, we proved that if M is a complete orientable noncompact hypersurface in $\mathbb{R}^{n+1}$ and δ-stability inequality holds on M, then M has at most one p-nonparabolic end. It was also proved that if M^n is a complete noncompact submanifold in $\mathbb{R}^{n+k}$ with sufficiently small L^n-norm of the traceless second fundamental form, then M has at most one p-nonparabolic end. For the reader's convenience, let us recall a definition of the p-nonparabolic ends. Let $E \subset M$ be an end of M, namely, E is an unbounded connected component of $M \setminus \Omega$ for a sufficiently large compact subset $\Omega \subset M$ with smooth boundary. As in usual harmonic function theory, we define the p-parabolicity and p-nonparabolicity of E as follows (see [Chang et al. 2016] and the references therein):

Definition 1.1. An end E of the Riemannian manifold M is called *p-parabolic* if for every compact subset $K \subset \overline{E}$

$$\mathrm{cap}_p(K, E) := \inf \int_E |\nabla f|^p = 0$$

where the infimum is taken among all $f \in C_0^\infty(\overline{E})$ such that $f \geq 1$ on K. Otherwise, the end E is called *p-nonparabolic*.

The first main result in this paper is the below theorem.

Theorem 1.2. *Let M^n $(n \geq 3)$ be a complete non-compact immersed submanifold of N^{n+m} with flat normal bundle. Assume that N^{n+m} has pure curvature tensor and the $(1, n-1)$-curvature of N^{n+m} is not less than $-k$. If one of the following conditions*

1. $$|A|^2 \leq \frac{n^2|H|^2 - 2k}{n-1},$$
2. $$\frac{n^2|H|^2 - 2k}{n-1} < |A|^2 \leq \frac{n^2|H|^2}{n-1} \text{ and } \lambda_1(M) > \frac{kp^2(n-1)}{4(p-1)(n+p-2)},$$
3. *the total curvature $\|A\|_n$ is bounded by*
$$\|A\|_n^2 < \min\left\{\frac{n^2}{(n-1)C_S}, \frac{2}{(n-1)C_S}\left[\frac{4(p-1)(n+p-2)}{p^2(n-1)} - \frac{k}{\lambda_1(M)}\right]\right\}$$
4. *$\sup_M |A|$ is bounded and the fundamental tone satisfies*
$$\lambda_1(M) > \frac{p^2(n-1)(2k + (n-1)\sup_M |A|^2)}{8(p-1)(n+p-2)},$$

holds true then every p-harmonic L^p 1-form on M is trivial. Therefore, M has at most one p-nonparabolic end. Here C_S is the Sobolev constant which depends only on n (see Lemma 2.1).

This results generalizes a work of the first author in [Dung 2017] On the other hand, we consider p-harmonic ℓ-forms on Riemannian manifolds with weighted Poincaré inequality. Recall that let (M^n, g) be a Riemannian manifold of dimension n and $\rho \in \mathcal{C}(M)$ be a positive function on M. We say that M has a weighted Poincaré inequality, if

$$\int_M \rho\varphi^2 \leq \int_M |\nabla\varphi|^2 \tag{1..1}$$

holds true for any smooth function $\varphi \in \mathcal{C}_0^\infty(M)$ with compact support in M. The positive function ρ is called the weighted function. It is easy to see that if the bottom of the spectrum of Laplacian $\lambda_1(M)$ is positive then M satisfies a weighted Poincaré inequality with $\rho \equiv \lambda_1$. Here $\lambda_1(M)$ can be characterized by variational principle

$$\lambda_1(M) = \inf\left\{\frac{\int_M |\nabla\varphi|^2}{\int_M \varphi^2} : \varphi \in \mathcal{C}_0^\infty(M)\right\}.$$

When M satisfies a weighted Poincaré inequality then M has many interesting properties concerning topology and geometry. For example, in [Vieira 2014], Vieira obtained vanishing theorems for L^2 harmonic 1 forms on complete Riemannian manifolds satisfying a weighted Poincaré inequality and having a certain lower bound of the bi-Ricci curvature. His theorems are improvement of Li-Wang's and Lam's results (see [Lam 2010, Li and Wang 2001, Li and Wang 2006]). Moreover, some applications to study geometric structures of minimal hypersurfaces also are given. We refer to [Chen and Sung 2009, Dung and Sung 2014] and the refrences therein for further results on the vanishing property of the space of harmonic ℓ-forms. In the nonlinear setting, Chang-Chen-Wei studied p-harmonic functions with finite L^q energy in [Chang et al. 2016], and proved some vanishing type theorems on Riemannian manifolds satisfying a weighted Poincaré inequality. Later, Sung-Wang, Dat and the first author used theory of p-harmonic functions to show some interesting rigidity propeties of Riemannian manifolds with maximal p-spectrum.

(see [Dung and Dat 2016, Sung and Wang 2014]). In this paper, we will investigate manifolds with weighted Poincaré inequality and prove some vanishing results for p-harmonic ℓ-forms on the manifolds. Our results can be considered as a generalization of Vieira's and the first author's results (see [Dung 2017]). Finally, we are also interested in locally conformally flat Riemannian manifolds, our theorem is the following vanishing property.

Theorem 1.3. *Let (M^n, g), $n \geq 3$, be an n-dimensional complete, simply connected, locally conformally flat Riemannian manifold. If one of the following conditions*

1. $$\|T\|_{n/2} + \frac{\|R\|_{n/2}}{\sqrt{n}} < \frac{4\left(p - 1 + \min\left\{1, \frac{(p-1)^2}{n-1}\right\}\right)}{Sp^2}.$$
2. *the scalar curvature R is nonpositive and*
$$K_{p,n} := \frac{p - 1 + \min\left\{1, \frac{(p-1)^2}{n-1}\right\}}{p^2} - \frac{n-1}{\sqrt{n}(n-2)} > 0$$
and
$$\|T\|_{n/2} < \frac{4K_{p,n}}{S} = 4K_{p,n}\mathcal{Y}(\mathbb{S}^n).$$

holds true, then every p-harmonic 1-form with finite $L^p (p \geq 2)$ norm on M is trivial, and M must has at most one p-nonparabolic end. Here T is the traceless Ricci tensor and S is the constant given in Lemma 5.1.

The rest of this paper is organized as follows. In Section 2, we recall some basic notations and useful fact on theory of smooth ℓ-forms. Then, we study p-harmonic ℓ-forms in submanifolds of N^{n+m} with flat normal bundle and pure curvature tensor. We will give a proof of Theorem 1.2 in this section. In Section 4, we derive some vanishing properties for p-harmonic ℓ-forms on manifolds with weighted Poincaré inequality. Finally, in Section 5, we consider locally conformally flat Riemannian manifolds and give a proof of Theorem 1.3.

2.. Preliminary notations

In this paper for $n \geq 3,\ 1 \leq \ell \leq n-1$ and $p \geq 2,\ q \geq 0$, we denote

$$C_{n,\ell} := \max\{\ell, n - \ell\}$$

and

$$A_{p,n,\ell} = \begin{cases} 1 + \dfrac{1}{\max \ell, n-\ell}, & \text{if } p = 2 \\ 1 + \dfrac{1}{(p-1)^2} \min\left\{1, \dfrac{(p-1)^2}{n-1}\right\}, & \text{if } p > 2 \text{ and } \ell = 1 \\ 1, & \text{if } p > 2 \text{ and } 1 < \ell \leq n-1. \end{cases}$$

Hence,

$$(p-1)^2(A_{p,n,\ell} - 1) = \begin{cases} \dfrac{1}{\max\{\ell, n-\ell\}}, & \text{if } p = 2 \text{ and } 1 \leq \ell \leq n-1 \\ \min\left\{1, \dfrac{(p-1)^2}{n-1}\right\}, & \text{if } p > 2 \text{ and } \ell = 1 \\ 0, & \text{if } p > 2 \text{ and } 1 < \ell \leq n-1. \end{cases}$$

We will use the following Sobolev inequality.

Lemma 2.1 ([Hoffman and Spruck 1974]). *Let M^n $(n \geq 3)$ be an n-dimensional complete submanifold in a complete simply-connected manifold with nonpositive sectional curvature. Then for any $f \in W_0^{1,2}(M)$ we have*

$$\left(\int_M |f|^{\frac{2n}{n-2}} dv\right)^{\frac{n-2}{n}} \leq C_S \int_M \left(|\nabla f|^2 + |H|^2 f^2\right) dv,$$

where C_S is the Sobolev constant which depends only on n.

An important ingredient in our methods is the following refined Kato inequality. In order to state the inequality, let us recall some notations. Suppose that $\{e_1, \ldots, e_n\}$ is an orthonormal basic of $\mathbb{R}^n$. Let $\theta_1 : \Lambda^{\ell+1}\mathbb{R}^n \to \mathbb{R} \otimes \Lambda^{\ell}\mathbb{R}^n$ given by

$$\theta_1(v_1 \wedge \ldots \wedge v_{\ell+1}) = \frac{1}{\sqrt{\ell+1}} \sum_{j=1}^{\ell+1} (-1)^j v_j \otimes v_1 \wedge \ldots \wedge \widehat{v_j} \wedge \ldots \wedge v_{\ell+1}$$

and $\theta_2 : \Lambda^{\ell-1}\mathbb{R}^n \to \mathbb{R} \otimes \Lambda^{\ell}\mathbb{R}^n$ by

$$\theta_2(\omega) = -\frac{1}{\sqrt{n+1-\ell}} \sum_{j=1}^{n} e_j \otimes (e_j \wedge \omega).$$

Lemma 2.2 ([Li 2012, Calderbank et al. 2000, Dung and Seo 2017]). *For $p \geq 2, \ell \geq 1$, let ω be an p-harmonic ℓ-form on a complete Riemannian manifold M^n. The following inequality holds*

$$\left|\nabla \left(|\omega|^{p-2}\omega\right)\right|^2 \geq A_{p,n,\ell} \left|\nabla |\omega|^{p-1}\right|^2. \tag{2..1}$$

Moreover, when $p = 2, \ell > 1$ then the equality holds if and only if there exixts a 1-form α such that

$$\nabla\omega = \alpha \otimes \omega - \frac{1}{\sqrt{\ell+1}}\theta_1(\alpha \wedge \omega) + \frac{1}{\sqrt{n+1-\ell}}\theta_2(i_\alpha \omega).$$

Proof. The inequality (2..1) is well-known when $\ell = 1, p = 2$, for example, see [Li 2012]. When $\ell = 1, p > 2$, the inequality (2..1) was proved by Seo and the first author in [Dung and Seo 2017]. Note that when $p = 2, \ell > 1$, the inequality (2..1) was proved by Calderbank et al. (see [Calderbank et al. 2000]) but we refer to [Cibotaru and Zhu 2012] in a way convenient for our purpose without introducing abstract notation.

Finally, when $p > 2, \ell > 1$, the inequality (2..1) is standard (see [Calderbank et al. 2000]). □

Note that if M is not complete, then (2..1) is still true provided that ω is a harmonic field, see [Cibotaru and Zhu 2012] for further discussion.

Suppose that M is a complete noncompact Riemannian manifold. Let $\{e_1, \ldots, e_n\}$ be a local orthonormal frame on M with dual coframe $\{\omega_1, \ldots, \omega_n\}$. Given an ℓ-form ω on M, the Weitzenböck curvature operator K_ℓ acting on ω is defined by

$$K_\ell = \sum_{j.k=1}^{n} \omega_k \wedge i_{e_j} R(e_k, e_j)\omega.$$

Using the Weitzenböck curvature operator, we have the following Bochner type formula for ℓ-forms.

Lemma 2.3 ([Li 2012]). *Let $\omega = \sum_I a_I \omega_I$ be a ℓ-form on M. Then*

$$\begin{aligned}\Delta|\omega|^2 &= 2\langle \Delta\omega, \omega\rangle + 2|\nabla\omega|^2 + 2\langle E(\omega), \omega\rangle \\ &= 2\langle \Delta\omega, \omega\rangle + 2|\nabla\omega|^2 + 2K_\ell(\omega, \omega)\end{aligned}$$

where $E(\omega) = \sum\limits_{j,k=1}^{n} \omega_k \wedge i_{e_j} R(e_k, e_j)\omega$.

In order to estimate the K_ℓ, we need to define a new curvature which is appear naturally as a component of the Weitzenböck curvature operator (see [Lin 2015a, Lin 2015c]).

Definition 2.4. Let M^n be a complete immersed submanifold in a Riemannian manifold N^{n+m} with flat normal bundle. Here the submanifold M is said to have flat normal bundle if the normal connection of M is flat, namely the components of the normal curvature tensor of M are zero. For any point $x \in N^{n+m}$, choose an orthonormal frame $\{e_i, \dots, e_n\}_{i=1}^{n+m}$ of the tangent space T_xN and define

$$\overline{R}^{(\ell,n-\ell)}([e_{i_1} \dots, e_{i_n}]) = \sum_{k=1}^{\ell} \sum_{h=\ell+1}^{n} \overline{R}_{i_k i_h i_k i_h}$$

for $1 \le \ell \le n-1$, where the indices $1 \le i_1, \dots i_n \le n+m$ are distinct with each other. We call $\overline{R}^{(\ell,n-\ell)}([e_{i_1} \dots, e_{i_n}])$ the $(\ell, n-\ell)$-curvature of N^{n+m}.

It is easy to see that if $\ell = 1$ then the $(1, n-1)$-curvature becomes $(n-1)$-th Ricci curvature ([Lin 2015c]). Assume that N has pure curvature tensor, namely for every $p \in N$ there is an orthonormal basis $\{e_1, \dots, e_n\}$ of the tangent space T_pN such that $R_{ijrs} := \langle R(e_i, e_j)e_r, e_s\rangle = 0$ whenever the set $\{i, j, r, s\}$ contains more than two elements. Here R_{ijrs} denote the curvature tensors of N. It is worth to notice that all 3-manifolds and conformally flat manifolds have pure curvature tensor. It was proved in [Lin 2015a] that

$$K_\ell(\omega, \omega) \ge \frac{1}{2}(n^2|H|^2 - C_{n,\ell}|A|^2) + \inf_{i_1,\dots,i_n} \overline{R}^{(\ell,n-\ell)}([e_{i_1} \dots, e_{i_n}])|\omega|^2.$$

Finally, to prove the vanishing property of p-harmonic ℓ-form, we use the following useful estimates. Since we can not find them in the literature, we will give a detail of discussion.

Lemma 2.5. *For any closed ℓ-form ω and $\varphi \in \mathcal{C}^\infty(M)$, we have*

$$|d(\varphi\omega)| = |d\varphi \wedge \omega| \le |d\varphi| \cdot |\omega|.$$

Proof. Let $\{X_i\}$ be a local orthonormal frame and $\{dx_i\}$ is the dual coframe. Since ω is closed, we have $d(\varphi\omega) = d\varphi \wedge \omega$. Suppose that

$$\omega = \sum_{|I|=\ell} \omega_I dx^I = \sum_{|K|=\ell-1} \sum_{j=1}^{n} \omega_{jK} dx_j \wedge dx^K,$$

where $\omega_{jK} = 0$ if $j \in K$. Therefore, denote $\varphi_i = \nabla_{X_i}\varphi$, we have

$$d\varphi \wedge \omega = \sum_{|K|=\ell-1} \sum_{i\ne j.i,j\notin K}^{n} \varphi_i \omega_{jK} dx_i \wedge dx_j \wedge dx^K.$$

Observe that, for any $a_i, b_i \in \mathbb{R}, i = \overline{1,n}$

$$\sum_{i<j} (a_i b_j - a_j b_i)^2 + \left(\sum_{i=1}^{n} a_i b_i\right)^2 = \left(\sum_{i=1}^{n} a_i^2\right)\left(\sum_{i=1}^{n} b_i^2\right),$$

we infer

$$\begin{aligned}
|d\varphi \wedge \omega|^2 &= \sum_{|K|=\ell-1} \sum_{i<j,i,j\notin K} (\varphi_i\omega_{jK} - \varphi_j\omega_{iK})^2 \\
&\le \sum_{|K|=\ell-1} \sum_{i<j} (\varphi_i\omega_{jK} - \varphi_j\omega_{iK})^2 \\
&\le \sum_{|K|=\ell-1} \left(\sum_{i=1}^{n} \varphi_i^2\right)\left(\sum_{j=1}^{n} \omega_{jK}\right) = |d\varphi|^2|\omega|^2.
\end{aligned}$$

The proof is complete. □

3.. p-HARMONIC ℓ-FORMS IN SUBMANIFOLDS OF N^{n+m} WITH FLAT NORMAL BUNDLE

Theorem 3.1. *Let M^n $(n \geq 3)$ be a complete non-compact immersed submanifold of N^{n+m} with flat normal bundle. Assume that N has pure curvature tensor and the $(\ell, n-\ell)$-curvature of N^{n+m} is not less than $-k$ for $1 \leq \ell \leq n-1$. If one of the following conditions*

1.
$$|A|^2 \leq \frac{n^2|H|^2 - 2k}{C_{n,\ell}}, \quad Vol(M) = \infty.$$
2.
$$\frac{n^2|H|^2 - 2k}{C_{n,\ell}} < |A|^2 \leq \frac{n^2|H|^2}{C_{n,\ell}} \text{ and } \lambda_1 > \frac{kQ^2}{4(Q-1+(p-1)^2(A_{p,n,\ell}-1))},$$
3. *the total curvature $\|A\|_n$ is bounded by*
$$\|A\|_n^2 < \min\left\{\frac{n^2}{C_{n,\ell}C_S}, \frac{2}{C_{n,\ell}C_S}\left[\frac{4(Q-1+(p-1)^2(A_{p,n,\ell}-1))}{Q^2} - \frac{k}{\lambda_1(M)}\right]\right\},$$
4. $\sup_M |A|$ *is bounded and the fundamental tone satisfies*
$$\lambda_1(M) > \frac{Q^2(2k + C_{n,\ell}\sup_M |A|^2)}{8(Q-1+(p-1)^2(A_{p,n,\ell}-1))},$$

holds true then every p-harmonic ℓ-form with finite L^Q, $(Q \geq 2)$ on M is trivial.

Proof. Let $M_+ := M \setminus \{x \in M, \omega(x) = 0\}$. Let ω be any p-harmonic ℓ-form with finite L^Q norm. Apply the Bochner formula to the form $|\omega|^{p-2}\omega$ we obtain, on M_+

$$\begin{aligned}\frac{1}{2}\Delta|\omega|^{2(p-1)} &= |\nabla(|\omega|^{p-2}\omega)|^2 - \left\langle(\delta d + d\delta)(|\omega|^{p-2}\omega), |\omega|^{p-2}\omega\right\rangle + K_\ell(|\omega|^{p-2}\omega, |\omega|^{p-2}\omega) \\ &= |\nabla(|\omega|^{p-2}\omega)|^2 - \left\langle\delta d(|\omega|^{p-2}\omega), |\omega|^{p-2}\omega\right\rangle + |\omega|^{2(p-2)}K_\ell(\omega,\omega)\end{aligned}$$

where we used ω is p-harmonic in the second equality. This can be read as

$$|\omega|^{p-1}\Delta|\omega|^{p-1} = \left(|\nabla(|\omega|^{p-2}\omega)|^2 - |\nabla|\omega|^{p-1}|^2\right) - |\omega|^{p-2}\left\langle\delta d(|\omega|^{p-2}\omega), \omega\right\rangle + |\omega|^{2(p-2)}K_\ell(\omega,\omega).$$

By Kato type inequality, we infer

$$|\omega|\Delta|\omega|^{p-1} \geq (p-1)^2(A_{p,n,\ell}-1)|\omega|^{p-2}|\nabla|\omega||^2 - \left\langle\delta d(|\omega|^{p-2}\omega), \omega\right\rangle + K_\ell|\omega|^p. \quad (3..1)$$

Hence,

$$|\omega|^{q+1}\Delta|\omega|^{p-1} \geq (p-1)^2(A_{p,n,\ell}-1)|\omega|^{p+q-2}|\nabla|\omega||^2 - \left\langle\delta d(|\omega|^{p-2}\omega), |\omega|^q\omega\right\rangle + K_\ell|\omega|^{p+q}.$$

We choose cutoff function $\varphi \in \mathcal{C}_0^\infty(M_+)$ then multiplying both sides of the above inequality by φ^2, we obtain

$$\begin{aligned}\int_{M_+}\varphi^2|\omega|^{q+1}\Delta|\omega|^{p-1} \geq &(p-1)^2(A_{p,n,\ell}-1)\int_{M_+}\phi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \\ &- \int_{M_+}\left\langle\delta d(|\omega|^{p-2}\omega), \varphi^2|\omega|^q\omega\right\rangle + \int_{M_+}K_\ell\varphi^2|\omega|^{p+q}.\end{aligned}$$

by integrating by part, this implies

$$\begin{aligned}\int_{M_+}\left\langle\nabla(\varphi^2|\omega|^{q+1}), \nabla|\omega|^{p-1}\right\rangle \leq &-(p-1)^2(A_{p,n,\ell}-1)\int_{M_+}\phi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \\ &+ \int_{M_+}\left\langle d(|\omega|^{p-2}\omega), d(\varphi^2|\omega|^q\omega)\right\rangle - \int_{M_+}K_\ell\varphi^2|\omega|^{p+q}.\end{aligned} \quad (3..2)$$

Since the $(\ell, n-\ell)$-curvature of N^{n+m} is not less than $-k$, we have

$$K_\ell \geq \frac{1}{2}(n^2|H|^2 - \max\{\ell, n-\ell\}|A|^2) - k.$$

Obviously,

$$\begin{aligned}\int_{M_+} \left\langle \nabla(\varphi^2|\omega|^{q+1}), \nabla|\omega|^{p-1}\right\rangle =& 2(p-1)\int_{M_+} \varphi|\omega|^{p+q-1}\left\langle \nabla\varphi, \nabla|\omega|\right\rangle \\ &+ (q+1)(p-1)\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2. \qquad (3..3)\end{aligned}$$

Note that for any close ℓ-form ω and smooth function f, it is proved in Lemma 2.5 that

$$d(f\wedge\omega) = df\wedge\omega \leq |df|.|\omega|.$$

Therefore,

$$\begin{aligned}\int_{M_+} \left\langle d(|\omega|^{p-2}\omega), d(\varphi^2|\omega|^q\omega)\right\rangle &= \int_{M_+} \left\langle d(|\omega|^{p-2})\wedge\omega, d(\varphi^2|\omega|^q)\wedge\omega\right\rangle \\ &\leq \int_{M_+} \left|d(|\omega|^{p-2})\wedge\omega|.|d(\varphi^2|\omega|^q)\wedge\omega\right| \\ &\leq \int_{M_+} \left|\nabla(|\omega|^{p-2})\right||\omega|.\left|\nabla(\varphi^2|\omega|^q)\right||\omega| \\ &\leq (p-2)q\int_{M_+} \varphi^2|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2 \\ &\quad + 2(p-2)\int_{M_+} \varphi|\omega|^{p+q-1}\left|\nabla|\omega|\right|\left|\nabla\varphi\right|. \qquad (3..4)\end{aligned}$$

1. If $|A|^2 \leq \dfrac{n^2|H|^2 - 2k}{C_{n,\ell}}$, then $K_\ell \geq 0$. Therefore, from (3..2) we obtain

$$\begin{aligned}\int_{M_+} \left\langle \nabla(\varphi^2|\omega|^{q+1}), \nabla|\omega|^{p-1}\right\rangle \leq& -(p-1)^2(A_{p,n,\ell}-1)\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \\ &+ \int_{M_+} \left\langle d(|\omega|^{p-2}\omega), d(\varphi^2|\omega|^q\omega)\right\rangle.\end{aligned}$$

Thus, by (3..3), (3..4),

$$\begin{aligned}&2(p-1)\int_{M_+} \varphi|\omega|^{p+q-1}\left\langle \nabla\varphi, \nabla|\omega|\right\rangle + (q+1)(p-1)\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \\ &\leq -(p-1)^2(A_{p,n,\ell}-1)\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \\ &\quad + (p-2)q\int_{M_+} \varphi^2|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2 + 2(p-2)\int_{M_+} \varphi|\omega|^{p+q-1}\left|\nabla|\omega|\right|\left|\nabla\varphi\right|.\end{aligned}$$

Hence,

$$(p+q-1+(p-1)^2(A_{p,n,\ell}-1))\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \leq 2(2p-3)\int_{M_+} \varphi|\omega|^{p+q-1}\left|\nabla|\omega|\right|\left|\nabla\varphi\right|.$$

Using the fundamental inequality $2AB \leq \varepsilon A^2 + \varepsilon^{-1}B^2$, we have that, for every $\varepsilon > 0$,

$$2\int_{M_+} \varphi|\omega|^{p+q-1}\left|\nabla|\omega|\right|\left|\nabla\varphi\right| \leq \varepsilon\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2 + \frac{1}{\varepsilon}\int_{M_+} |\omega|^{p+q}|\nabla\varphi|^2. \qquad (3..5)$$

From the last two inequalities, we obtain

$$(p+q-1+(p-1)^2(A_{p,n,\ell}-1)-\varepsilon(2p-3))\int_{M_+} \varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2 \leq \frac{2p-3}{\varepsilon}\int_{M_+} |\omega|^{p+q}|\nabla\varphi|^2.$$

Let $Q := p+q$, since $Q-1+(p-1)^2(A_{p,n,\ell}-1) > 0$, we can choose $\varepsilon > 0$ small enough and a constant $K = K(\varepsilon) > 0$ so that

$$\frac{4}{Q^2}\int_{M_+}\varphi^2\left|\nabla|\omega|^{Q/2}\right|^2 \leq K\int_{M_+}|\omega|^Q|\nabla\varphi|^2, \quad \text{for all } r>0. \tag{3..6}$$

Applying a variation of the Duzaar-Fuchs cut-off method (see also [Duzaar and Fuchs 1990, Nakauchi 1998]), we shall show that (3..6) holds for every $\varphi \in C_0^\infty(M)$. Indeed, we define

$$\eta_{\tilde\epsilon} = \min\left\{\frac{|\omega|}{\tilde\epsilon}, 1\right\}$$

for $\tilde\epsilon > 0$. Let $\varphi_{\tilde\epsilon} = \psi^2\eta_{\tilde\epsilon}$, where $\psi \in \mathcal{C}_0^\infty(M)$. It is easy to see that $\varphi_{\tilde\epsilon}$ is a compactly supported continuous function and $\varphi_{\tilde\epsilon} = 0$ on $M\setminus M_+$. Now, we replace φ by $\varphi_{\tilde\epsilon}$ in (3..6) and get

$$\begin{aligned}&\int_{M_+}\psi^4(\eta_{\tilde\epsilon})^2|\omega|^{Q-2}|\nabla|\omega||^2\\ &\qquad\leq 6C\int_{M_+}|\omega|^Q|\nabla\psi|^2\psi^2(\eta_{\tilde\epsilon})^2 + 3C\int_{M_+}|\omega|^Q|\nabla\eta_{\tilde\epsilon}|^2\psi^4.\end{aligned} \tag{3..7}$$

Observe that

$$\int_{M_+}|\omega|^Q|\nabla\eta_{\tilde\epsilon}|^2\psi^4 \leq \tilde\epsilon^{Q-2}\int_{M_+}|\nabla|\omega||^2\psi^4\chi_{\{|\omega|\leq\tilde\epsilon\}}$$

and the right hand side vanishes by dominated convergence as $\tilde\epsilon\to 0$, because $|\nabla|\omega|| \in L^2_{loc}(M)$ and $Q\geq 2$. Letting $\tilde\epsilon\to 0$ and applying Fatou lemma to the integral on the left hand side and dominated convergence to the first integral in the right hand side of (3..7), we obtain

$$\int_{M_+}\psi^4|\omega|^{Q-2}|\nabla|\omega||^2 \leq 6C\int_{M_+}|\omega|^Q|\nabla\psi|^2\psi^2, \tag{3..8}$$

where $\psi\in C_0^\infty(M)$. We choose cutoff functions $\psi\in\mathcal{C}_0^\infty(M_+)$ satisfying

$$\psi = \begin{cases}1, & \text{on } B_r\\ \in[0,1] \text{ and } |\nabla\varphi|\leq\frac{2}{r}, & \text{on } B_{2r}\setminus B_r\\ 0, & \text{on } M\setminus B_{2r}\end{cases},$$

where B_r is the open ball of radius r and center at a fixed point of M.

Letting $r\to\infty$, we conclude that $|\omega|$ is constant on M_+. Since $|\omega| = 0 \in \partial M_+$, it implies that either ω is zero or $M_+ = \emptyset$. If $M_+ = \emptyset$, then $|\omega|$ is contant on M. Thanks to the assumption $|\omega|\in L^Q(M)$, we infer $\omega = 0$.

2. Assume that

$$\frac{n^2|H|^2-2k}{C_{n,\ell}} < |A|^2 \leq \frac{n^2|H|^2}{C_{n,\ell}}$$

then $-k\leq K_\ell < 0$. From this and (3..2) we obtain

$$\begin{aligned}\int_{M_+}\left\langle\nabla(\varphi^2|\omega|^{q+1}),\nabla|\omega|^{p-1}\right\rangle \leq &-(p-1)^2(A_{p,n,\ell}-1)\int_{M_+}|\omega|^{p+q-2}|\nabla|\omega||^2\\ &\int_{M_+}\left\langle d(|\omega|^{p-2}\omega), d(\varphi^2|\omega|^q\omega)\right\rangle + k\int_{M_+}\varphi^2|\omega|^{p+q}.\end{aligned} \tag{3..9}$$

By the definition of $\lambda_1(M)$ and (3..5), we obtain that, for any $\varepsilon>0$,

$$\begin{aligned}\lambda_1\int_{M_+}\varphi^2|\omega|^{p+q} \leq &\int_{M_+}\left|\nabla\left(\varphi|\omega|^{(p+q)/2}\right)\right|^2\\ =&\frac{(p+q)^2}{4}\int_{M_+}\varphi^2|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2 + \int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2\end{aligned}$$

$$+ (p+q)\int_{M_+} \varphi|\omega|^{p+q-1}\langle\nabla|\omega|,\nabla\varphi\rangle$$
$$\leq (1+\varepsilon)\frac{(p+q)^2}{4}\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2+\left(1+\frac{1}{\varepsilon}\right)\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2. \tag{3..10}$$

From this, (3..3), (3..4), and (3..9), we have

$$C_\varepsilon\int_{M_+}\varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2\leq D_\varepsilon\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2,$$

where

$$C_\varepsilon := p+q-1+(p-1)^2(A_{p,n,\ell}-1)-\varepsilon(2p-3)-(1+\varepsilon)\frac{k(p+q)^2}{4\lambda_1(M)}$$
$$= Q-1+(p-1)^2(A_{p,n,\ell}-1)-\varepsilon(2p-3)-(1+\varepsilon)\frac{kQ^2}{4\lambda_1(M)},$$

and

$$D_\varepsilon := \frac{2p-3}{\varepsilon}+\frac{k}{\lambda_1(M)}\left(1+\frac{1}{\varepsilon}\right).$$

Here $Q=p+q\geq 2$. Since

$$\lambda_1(M)>\frac{kQ^2}{4(Q-1+(p-1)^2(A_{p,n,\ell}-1))},$$

there are some small enough number $\varepsilon>0$ and constant $K=K(\varepsilon)>0$ such that

$$\frac{4}{Q^2}\int_{M_+}\left|\nabla|\omega|^{Q/2}\right|^2\leq K\int_{M_+}|\omega|^Q|\nabla\varphi|^2.$$

Arguing similarly as in the proof of the first part, we conclude that $|\omega|$ is constant. Since $\lambda_1(M)>0$, M must have infinite volume, note that $|\omega|\in L^Q(M)$, we have that ω is zero.

3. Assume that $|A|^2>\dfrac{n^2|H|^2}{C_{n,\ell}}$. Then, from (3..2) we have

$$\int_{M_+}\left\langle\nabla(\varphi^2|\omega|^{q+1}),\nabla|\omega|^{p-1}\right\rangle+\frac{n^2}{2}\int_{M_+}\varphi^2|H|^2|\omega|^{p+q}$$
$$\leq -(p-1)^2(A_{p,n,\ell}-1)\int_{M_+}\varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2$$
$$+\int_{M_+}\left\langle d(|\omega|^{p-2}\omega),d(\varphi^2|\omega|^q\omega)\right\rangle+\frac{C_{n,\ell}}{2}\int_{M_+}\varphi^2|A|^2|\omega|^{p+q}+k\int_{M_+}\varphi^2|\omega|^{p+q}. \tag{3..11}$$

By Hölder inequality and Lemma 2.1, we have

$$\int_{M_+}\varphi^2|A|^2|\omega|^{p+q}\leq\|A\|_n^2\left(\int_{M_+}\left(\varphi|\omega|^{(p+q)/2}\right)^{\frac{2n}{n-2}}\right)^{\frac{n-2}{n}}$$
$$\leq C_S\|A\|_n^2\left(\int_{M_+}\left|\nabla\left(\varphi|\omega|^{(p+q)/2}\right)\right|^2+\int_{M_+}\varphi^2|H|^2|\omega|^{p+q}\right),$$

where C_S is the Sobolev constant depending only on n. From the last inequality and (3..5) we can get that, for any $\varepsilon>0$,

$$\int_{M_+}\varphi^2|A|^2|\omega|^{p+q}\leq C_S\|A\|_n^2(1+\varepsilon)\frac{(p+q)^2}{4}\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2$$
$$+C_S\|A\|_n^2\left(1+\frac{1}{\varepsilon}\right)\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2+C_S\|A\|_n^2\int_{M_+}\varphi^2|H|^2|\omega|^{p+q}.$$

Using this inequality and (3..3), (3..4), (3..11), we have

$$C_\varepsilon \int_{M_+} \varphi^2 |\omega|^{p+q-2} |\nabla |\omega||^2 + \frac{1}{2}\left(n^2 - C_{n,\ell} C_S \|A\|_n^2\right) \int_{M_+} \varphi^2 |H|^2 |\omega|^{p+q} \le D_\varepsilon \int_{M_+} |\omega|^{p+q} |\nabla \varphi|^2,$$

where, for $Q := p+q \ge 2$

$$\begin{aligned} C_\varepsilon :&= p+q-1+(p-1)^2(A_{p,n,\ell}-1) - \varepsilon(2p-3) - (1+\varepsilon)\frac{(p+q)^2}{4}\left(\frac{k}{\lambda_1(M)} + \frac{C_{n,\ell} C_S \|A\|_n^2}{2}\right) \\ &= Q-1+(p-1)^2(A_{p,n,\ell}-1) - \varepsilon(2p-3) - (1+\varepsilon)\frac{Q^2}{4}\left(\frac{k}{\lambda_1(M)} + \frac{C_{n,\ell} C_S \|A\|_n^2}{2}\right), \end{aligned}$$

and

$$D_\varepsilon := \frac{2p-3}{\varepsilon} + \left(1+\frac{1}{\varepsilon}\right)\left(\frac{k}{\lambda_1(M)} + \frac{C_{n,\ell} C_S \|A\|_n^2}{2}\right).$$

Since

$$\|A\|_n^2 < \min\left\{\frac{n^2}{C_{n,\ell} C_S},\ \frac{2}{C_{n,\ell} C_S}\left[\frac{4(Q-1+(p-1)^2(A_{p,n,\ell}-1))}{Q^2} - \frac{k}{\lambda_1(M)}\right]\right\},$$

there are some small enough $\varepsilon > 0$ and constant $K = K(\varepsilon) > 0$ such that

$$\frac{4}{Q^2}\int_{M_+} \left|\nabla |\omega|^{Q/2}\right|^2 \varphi^2 \le K \int_{M_+} |\omega|^Q |\nabla \varphi|^2.$$

Using the same argument as in the proof of the first and second part, this inequality implies that ω is zero.

4. Suppose that $\sup_M |A|^2 < \infty$. Then using (3..10) we have that, for any $\varepsilon > 0$,

$$\begin{aligned} \int_{M_+} \varphi^2 |A|^2 |\omega|^{p+q} &\le \sup_M |A|^2 \int_{M_+} \varphi^2 |\omega|^{p+q} \le \frac{\sup_M |A|^2}{\lambda_1(M)} \int_{M_+} \left|\nabla\left(\varphi |\omega|^{(p+q)/2}\right)\right|^2 \\ &\le \frac{\sup_M |A|^2}{\lambda_1(M)}(1+\varepsilon)\frac{(p+q)^2}{4}\int_{M_+} \varphi^2 |\omega|^{p+q-2} |\nabla |\omega||^2 \\ &\quad + \frac{\sup_M |A|^2}{\lambda_1(M)}\left(1+\frac{1}{\varepsilon}\right)\int_{M_+} |\omega|^{p+q} |\nabla \varphi|^2. \end{aligned}$$

From this and (3..3), (3..4), (3..11) we obtain that, for any $\varepsilon > 0$,

$$C_\varepsilon \int_{M_+} \varphi^2 |\omega|^{p+q-2} |\nabla |\omega||^2 + \frac{n^2}{2}\int_{M_+} \varphi^2 |H|^2 |\omega|^{p+q} \le D_\varepsilon \int_{M_+} |\omega|^{p+q} |\nabla \varphi|^2,$$

where

$$\begin{aligned} C_\varepsilon :&= p+q-1+(p-1)^2(A_{p,n,\ell}-1) - \varepsilon(2p-3) - (1+\varepsilon)\frac{(p+q)^2}{4\lambda_1(M)}\left(k + \frac{C_{n,\ell} \sup_M |A|^2}{2}\right) \\ &= Q-1+(p-1)^2(A_{p,n,\ell}-1) - \varepsilon(2p-3) - (1+\varepsilon)\frac{Q^2}{4\lambda_1(M)}\left(k + \frac{C_{n,\ell} \sup_M |A|^2}{2}\right), \end{aligned}$$

and

$$D_\varepsilon := \frac{2p-3}{\varepsilon} + \left(1+\frac{1}{\varepsilon}\right)\frac{1}{\lambda_1(M)}\left(k + \frac{C_{n,\ell} \sup_M |A|^2}{2}\right).$$

Since

$$\lambda_1(M) > \frac{Q^2(2k + C_{n,\ell} \sup_M |A|^2)}{8(Q-1+(p-1)^2(A_{p,n,\ell}-1))},$$

M must have infinite volume. Moreover, there are some small enough $\varepsilon > 0$ and constant $K = K(\varepsilon) > 0$ such that

$$\frac{4}{Q^2}\int_{M_+} \left|\nabla |\omega|^{Q/2}\right|^2 \varphi^2 \le K \int_{M_+} |\omega|^Q |\nabla \varphi|^2.$$

Using the same argument as in the proof of the first and second part, this inequality implies that ω is zero. □

Now, we will give a geometric application of Theorem 3.1 that is the Theorem 1.2. First, let us recall the following result about the existence of p-harmonic function on a Riemannian manifold.

Theorem 3.2 ([Chang et al. 2016]). *Let M be a Riemannian manifold with at least two p-nonparabolic ends. Then, there exists a non-constant, bounded p-harmonic function $u \in \mathcal{C}^{1,\alpha}(M)$ for some $\alpha > 0$ such that $|\nabla u| \in L^p(M)$.*

Note that, it is known that the regularity of (weakly) p-harmonic function u is not better than $\mathcal{C}^{1,\alpha}_{loc}$ (see [Tolksdorff 1984] and the references therein). Moreover $u \in W^{2,2}_{loc}$ if $p \geq 2$; $u \in W^{2,p}_{loc}$ if $1 < p < 2$ (see [Tolksdorff 1984]). In fact, any nontrivial (weakly) p-harmonic function u on M is smooth away from the set $\{\nabla u = 0\}$ which has n-dimensional Hausdorff measure zero.

Proof of Theorem 1.2. The proof follows by combining Theorem 3.1 with $q = 0, l = 1$ and Theorem 3.2. □

4.. p-HARMONIC ℓ-FORMS ON RIEMANNIAN MANIFOLDS WITH WEIGHTED POINCARÉ INEQUALITY

Lemma 4.1. *Let M be a complete Riemannian manifold satisfying a weighted Poincaré inequality with a continuous weight function ρ. Suppose that ω is a closed ℓ-form with finite L^{p+q} norm on M satisfies the following differential inequality*

$$|\omega|\Delta|\omega|^{p-1} \geq B|\omega|^{p-2}\left|\nabla|\omega|\right|^2 - \left\langle \delta d(|\omega|^{p-2}\omega), \omega\right\rangle - a\rho|\omega|^p - b|\omega|^p, \tag{4..1}$$

for some constants $0 < a < \dfrac{4(Q-1+B)}{Q^2}, b > 0$ and $Q \geq 2$. Then the following integral inequality holds

$$\int_{M_+}\left|\nabla|\omega|^{Q/2}\right|^2 \leq \frac{bQ^2}{4(Q-1+B)-aQ^2}\int_{M_+}|\omega|^Q. \tag{4..2}$$

Moreover, if equality holds in (4..2), then equality holds in (4..1)

Proof. (i) Assume that the manifold is compact. Multiplying inequality (4..1) by $|\omega|^q$ and then integrating by parts, we obtain

$$\begin{aligned}\int_{M_+}|\omega|^{q+1}\Delta|\omega|^{p-1} \geq & B\int_{M_+}|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2 - \int_{M_+}\left\langle \delta d(|\omega|^{p-2}\omega), |\omega|^q\omega\right\rangle \\ & - a\int_{M_+}\rho|\omega|^{p+q} - b\int_{M_+}|\omega|^{p+q},\end{aligned}$$

and then,

$$\begin{aligned}[(q+1)(p-1)+B]\int_{M_+}|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2 \leq & \int_{M_+}\left\langle d(|\omega|^{p-2}\omega), d(|\omega|^q\omega)\right\rangle \\ & + a\int_{M_+}\rho|\omega|^{p+q} + b\int_{M_+}|\omega|^{p+q}.\end{aligned}$$

Similarly to (3..4), we have

$$\int_{M_+}\left\langle d(|\omega|^{p-2}\omega), d(|\omega|^q\omega)\right\rangle = \int_{M_+}\left\langle d(|\omega|^{p-2})\wedge\omega, d(|\omega|^q)\wedge\omega\right\rangle$$

$$\begin{aligned}
&\leq \int_{M_+} \left|d(|\omega|^{p-2})\wedge\omega|.|d(|\omega|^q)\wedge\omega\right| \\
&\leq \int_{M_+} \left|\nabla(|\omega|^{p-2})\right||\omega|.\left|\nabla(|\omega|^q)\right||\omega| \\
&= (p-2)q\int_{M_+}|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2.
\end{aligned}$$

By the weighted Poincaré inequality we have that

$$\int_{M_+}\rho|\omega|^{p+q}\leq\int_{M_+}\left|\nabla\left(|\omega|^{(p+q)/2}\right)\right|^2=\frac{(p+q)^2}{4}\int_{M_+}|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2.$$

Combining the last three inequalities, we obtain

$$\left[p+q-1+B-a\frac{(p+q)^2}{4}\right]\int_{M_+}|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2\leq b\int_{M_+}|\omega|^{p+q},$$

consequently,

$$\left[\frac{4(p+q-1+B)}{(p+q)^2}-a\right]\int_{M_+}\left|\nabla|\omega|^{(p+q)/2}\right|^2\leq b\int_{M_+}|\omega|^{p+q}.$$

Let $Q:=p+q$, the last inequality implies that inequality (4..2) holds.

Now assume that equality holds in (4..2). Multiplying inequality (4..1) by $|\omega|^q$ where $q=Q-p$, and then integrating by parts and using the above estimates, we obtain

$$\begin{aligned}
0&\leq\int_{M_+}\left(|\omega|^{q+1}\Delta|\omega|^{p-1}-B|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2+\left\langle\delta d(|\omega|^{p-2}\omega),|\omega|^q\omega\right\rangle+a\rho|\omega|^{p+q}+b|\omega|^{p+q}\right)\\
&=-[(q+1)(p-1)+B]\int_{M_+}|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2+\int_{M_+}\left\langle d(|\omega|^{p-2}\omega),d(|\omega|^q\omega)\right\rangle\\
&+a\int_{M_+}\rho|\omega|^{p+q}+b\int_{M_+}|\omega|^{p+q}\\
&\leq-\left[\frac{4(p+q-1+B)}{(p+q)^2}-a\right]\int_{M_+}\left|\nabla|\omega|^{(p+q)/2}\right|^2+b\int_{M_+}|\omega|^{p+q}\\
&=0.
\end{aligned}$$

Therefore, we can conclude that equality holds in (4..1) in M_+. Since in $M\setminus M_+$, (4..1) is always true. We complete the proof.

(ii) Assume that the manifold is non-compact. We choose cutoff function $\varphi\in\mathcal{C}_0^\infty(M_+)$ as in the proof of Theorem 3.1. Multiplying both sides of inequality (4..1) by $\varphi^2|\omega|^q$ and then integrating by parts, we obtain

$$\begin{aligned}
\int_{M_+}\varphi^2|\omega|^{q+1}\Delta|\omega|^{p-1}\geq&B\int_{M_+}\varphi^2|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2-\int_{M_+}\left\langle\delta d(|\omega|^{p-2}\omega),\varphi^2|\omega|^q\omega\right\rangle\\
&-a\int_{M_+}\rho\varphi^2|\omega|^{p+q}-b\int_{M_+}\varphi^2|\omega|^{p+q},
\end{aligned}$$

and then,

$$\begin{aligned}
&\int_{M_+}\left\langle\nabla(\varphi^2|\omega|^{q+1}),\nabla|\omega|^{p-1}\right\rangle+B\int_{M_+}\varphi^2|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2\\
&\leq\int_{M_+}\left\langle d(|\omega|^{p-2}\omega),d(\varphi^2|\omega|^q\omega)\right\rangle+a\int_{M_+}\rho\varphi^2|\omega|^{p+q}+b\int_{M_+}\varphi^2|\omega|^{p+q},
\end{aligned}$$

Using the last inequality and (3..3), (3..4), we have

$$(p+q-1+B)\int_{M_+}\varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2\leq a\int_{M_+}\rho\varphi^2|\omega|^{p+q}+b\int_{M_+}\varphi^2|\omega|^{p+q}$$

$$+ 2(2p-3)\int_{M_+} \varphi|\omega|^{p+q-1}\,|\nabla|\omega||\,|\nabla\varphi|.$$

Similarly to (3..10) and by the weighted Poincaré inequality we have that

$$\begin{aligned}\int_{M_+} \rho\varphi^2|\omega|^{p+q} &\le \int_{M_+} \left|\nabla\left(\varphi|\omega|^{(p+q)/2}\right)\right|^2\\ &\le (1+\varepsilon)\frac{(p+q)^2}{4}\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2 + \left(1+\frac{1}{\varepsilon}\right)\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2.\end{aligned}$$

Combining the last two inequalities and using (3..5), we obtain

$$\begin{aligned}&\left[p+q-1+B-\varepsilon(2p-3)-(1+\varepsilon)\frac{a(p+q)^2}{4}\right]\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2\\ &\le b\int_{M_+}\varphi^2|\omega|^{p+q} + \left[\frac{2p-3}{\varepsilon}+a\left(1+\frac{1}{\varepsilon}\right)\right]\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2.\end{aligned}$$

Since $4(p+q-1+B)-a(p+q)^2>0$, $p+q-1+B-(1+\varepsilon)\dfrac{a(p+q)^2}{4}-\varepsilon(2p-3)>0$ for all sufficiently small enough $\varepsilon>0$. By the monotone convergence theorem, letting $r\to\infty$, and then $\varepsilon\to 0$, we obtain inequality (4..2).

Now suppose that equality in (4..2) holds. Multiplying both sides of inequality (4..1) by $\varphi^2|\omega|^q$ and then integrating by parts and using the above estimates, we obtain

$$\begin{aligned}0 &\le \int_{M_+}\varphi^2\left(|\omega|^{q+1}\Delta|\omega|^{p-1} - B|\omega|^{p+q-2}\,|\nabla|\omega||^2 + \left\langle\delta d(|\omega|^{p-2}\omega),|\omega|^q\omega\right\rangle + a\rho|\omega|^{p+q}+b|\omega|^{p+q}\right)\\ &\le -\left[p+q-1+B\right]\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2 + 2(2p-3)\int_{M_+}\varphi|\omega|^{p+q-1}|\nabla|\omega||\,|\nabla\varphi|\\ &\quad + a\int_{M_+}\rho\varphi^2|\omega|^{p+q} + b\int_{M_+}\varphi^2|\omega|^{p+q}\\ &\le -\left[p+q-1+B-\varepsilon(2p-3)-(1+\varepsilon)\frac{a(p+q)^2}{4}\right]\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2\\ &\quad + b\int_{M_+}\varphi^2|\omega|^{p+q} + \left[\frac{2p-3}{\varepsilon}+a\left(1+\frac{1}{\varepsilon}\right)\right]\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2.\end{aligned}$$

Letting $r\to\infty$ in the last inequality and using the monotone convergence theorem, we get

$$\begin{aligned}0 &\le \int_{M_+}\left(|\omega|^{q+1}\Delta|\omega|^{p-1} - B|\omega|^{p+q-2}\,|\nabla|\omega||^2 + \left\langle\delta d(|\omega|^{p-2}\omega),|\omega|^q\omega\right\rangle + a\rho|\omega|^{p+q}+b|\omega|^{p+q}\right)\\ &\le -\left[p+q-1+B-\varepsilon(2p-3)-(1+\varepsilon)\frac{a(p+q)^2}{4}\right]\int_{M_+}|\omega|^{p+q-2}\,|\nabla|\omega||^2 + b\int_{M_+}|\omega|^{p+q}.\end{aligned}$$

And then putting $\varepsilon\to 0$, we obtain

$$\begin{aligned}0 &\le \int_{M_+}\left(|\omega|^{q+1}\Delta|\omega|^{p-1} - B|\omega|^{p+q-2}\,|\nabla|\omega||^2 + \left\langle\delta d(|\omega|^{p-2}\omega),|\omega|^q\omega\right\rangle + a\rho|\omega|^{p+q}+b|\omega|^{p+q}\right)\\ &\le -\left[p+q-1+B-\frac{a(p+q)^2}{4}\right]\int_{M_+}|\omega|^{p+q-2}\,|\nabla|\omega||^2 + b\int_{M_+}|\omega|^{p+q}\\ &= 0.\end{aligned}$$

In view of (4..1), we can conclude that equality holds in (4..1). □

The following result was showed by Vieira in [Vieira 2014].

Lemma 4.2 ([Vieira 2014])**.** *Suppose that u is a smooth function on a complete Riemannian manifold M with finite L^Q norm, for $Q \geq 2$. Then*

$$\lambda_1(M) \int_M |u|^Q \leq \int_M \left|\nabla u^{Q/2}\right|^2.$$

Theorem 4.3. *Let M^n be a complete non-compact Riemannian manifold M^n satisfying a weighted Poincaré inequality with a continuous weight function ρ. Suppose that the curvature operator acting on ℓ-forms has a lower bound*

$$K_\ell \geq -a\rho - b$$

for some constants $b > 0, Q \geq 2$ and

$$0 < a < \frac{4(Q - 1 + (p-1)^2(A_{p,n,\ell} - 1))}{Q^2}.$$

Assume that the first eigenvalue of the Laplacian has a lower bound

$$\lambda_1(M^n) > \frac{bQ^2}{4(Q - 1 + (p-1)^2(A_{p,n,\ell} - 1)) - aQ^2}.$$

Then the space of p-harmonic ℓ-forms with finite L^Q energy on M^n is trivial.

Proof. Let ω be any p-harmonic ℓ-form with finite L^Q norm. From the Bochner formula and the Kato type inequality (see, Lemma 2.2), we obtain (3..1)

$$|\omega|\Delta|\omega|^{p-1} \geq (p-1)^2(A_{p,n,\ell} - 1)|\omega|^{p-2}\left|\nabla|\omega|\right|^2 - \left\langle \delta d(|\omega|^{p-2}\omega), \omega \right\rangle + K_\ell|\omega|^p.$$

Hence,

$$|\omega|\Delta|\omega|^{p-1} \geq (p-1)^2(A_{p,n,\ell} - 1)|\omega|^{p-2}\left|\nabla|\omega|\right|^2 - \left\langle \delta d(|\omega|^{p-2}\omega), \omega \right\rangle - a\rho|\omega|^p - b|\omega|^p.$$

Applying Lemma 4.1 to $B = (p-1)^2(A_{p,n,\ell} - 1)$, we obtain

$$\int_{M_+} \left|\nabla|\omega|^{Q/2}\right|^2 \leq \frac{bQ^2}{4(Q - 1 + (p-1)^2(A_{p,n,\ell} - 1)) - aQ^2} \int_{M_+} |\omega|^Q.$$

Since $|\omega| \in L^Q(M)$ this implies $|\omega| \in L^Q(M_+)$. Therefore, by Lemma 4.2, we have

$$\lambda_1(M^n) \int_{M_+} |\omega|^Q \leq \int_{M_+} \left|\nabla\omega^{Q/2}\right|^2.$$

From the last two inequalities, we obtain

$$\lambda_1(M^n) \int_{M_+} |\omega|^Q \leq \frac{bQ^2}{4(Q - 1 + (p-1)^2(A_{p,n,\ell} - 1)) - aQ^2} \int_{M_+} |\omega|^Q.$$

If the ℓ-form ω is not identically zero in M_+ (therefore $\omega = 0$ in M), then

$$\lambda_1(M^n) \leq \frac{bQ^2}{4(Q - 1 + (p-1)^2(A_{p,n,\ell} - 1)) - aQ^2},$$

which leads to a contradiction. So ω is identically zero. □

Combining Theorem 3.2 and Theorem 4.3 with $Q = p \geq 2, \ell = 1$, we obtain the follows result.

Corollary 4.4. *Let M^n be a complete non-compact Riemannian manifold M^n satisfying a weighted Poincaré inequality with a continuous weight function ρ. Suppose that*

$$Ric_M \geq -a\rho - b, \quad \text{for } 0 < a < \frac{4(p-1)(p+n-2)}{p^2(n-1)}$$

and some constants $b > 0$. Assume that the first eigenvalue of the Laplacian has a lower bound

$$\lambda_1(M^n) > \frac{bp^2(n-1)}{4(p-1)(n+p-2) - ap^2(n-1)}.$$

Then the space of L^p p-harmonic 1-forms on M^n is trivial. Therefore, M has at most one p-nonparabolic end.

Let

$$A_{n,\ell} = \begin{cases} \dfrac{n-\ell+1}{n-\ell}, & \text{if } 1 \le \ell \le \frac{n}{2} \\ \dfrac{\ell+1}{\ell}, & \text{if } \frac{n}{2} \le \ell \le n-1. \end{cases}$$

We conclude this section by the below rigidity property.

Corollary 4.5. *Let M^n be a complete non-compact Riemannian manifold M^n satisfying a weighted Poincaré inequality with a continuous weight function ρ. Suppose that*

$$Ric_M \ge -a\rho - b, \quad \text{for } 0 < a < A_{n,\ell}$$

and some constants $b > 0$. Assume that

$$\lambda_1(M^n) = \frac{b}{A_{n,\ell} - a}.$$

Then either

(1) *The space of L^2 harmonic ℓ-forms on M^n is trivial or;*
(2) *For any L^2 harmonic ℓ-form ω on M, there exixts a 1-form α such that*

$$\nabla\omega = \alpha \otimes \omega - \frac{1}{\sqrt{\ell+1}}\theta_1(\alpha \wedge \omega) + \frac{1}{\sqrt{n+1-\ell}}\theta_2(i_\alpha \omega).$$

Proof. Let $p = 2, q = 0, Q = 2$, the Bochner formula applying on harmonic ℓ-form ω implies that

$$|\omega|\Delta|\omega| \ge (A_{n,\ell} - 1)|\nabla|\omega||^2 - a\rho|\omega|^2 - b|\omega|^2.$$

If ω is non-trivial then Lemma 4.1 and Lemma 4.2 imply that

$$\lambda_1(M)\int_{M_+} |\omega|^2 = \int_{M_+} |\nabla|\omega||^2.$$

Therefore,

$$|\omega|\Delta|\omega| = (A_{n,\ell} - 1)|\nabla|\omega||^2 - a\rho|\omega|^2 - b|\omega|^2$$

on M_+, hence it holds true on M. This means that the equality in the Kato type inequality (2..1) holds true. By Lemma 2.2, we are done. □

5.. p-HARMONIC 1-FORMS ON LOCALLY CONFORMALLY FLAT RIEMANNIAN MANIFOLDS

In this section we prove vanishing theorem for p-harmonic 1-forms ($Q \ge 2$) with finite L^Q energy. We will use the following auxiliary lemmas.

It is known that a simply connected, locally conformally flat manifold M^n, $n \ge 3$, has a conformal immersion into $\mathbb{S}^n$, and according to [Schoen and Yau 1994], the Yamabe constant of M^n satisfies

$$\mathcal{Y}(M^n) = \mathcal{Y}(\mathbb{S}^n) = \frac{n(n-2)\omega_n^{2/n}}{4},$$

where ω_n is the volume of the unit sphere in $\mathbb{R}^n$. Therefore the following inequality

$$\mathcal{Y}(\mathbb{S}^n)\left(\int_M f^{\frac{2n}{n-2}}dv\right)^{\frac{n-2}{n}} \le \int_M |\nabla f|^2 dv + \frac{n-2}{4(n-1)}\int_M Rf^2 dv \qquad (5..1)$$

holds for all $f \in C_0^\infty(M)$. To derive vanishing property of p-harmonic 1-forms, we need to have following lemmas.

Lemma 5.1. *([Lin 2015b]) Let (M^n, g), $n \geq 3$, be an n-dimensional complete, simply connected, locally conformally flat Riemannian manifold with $R \leq 0$ or $\|R\|_{n/2} < \infty$. Then the following L^2 Sobolev inequality*

$$\left(\int_M f^{\frac{2n}{n-2}} dv\right)^{\frac{n-2}{n}} \leq S \int_M |\nabla f|^2 dv, \quad \forall f \in C_0^\infty(M),$$

holds for some constant $S > 0$, which is equal to $\mathcal{Y}(\mathbb{S}^n)^{-1}$ in the case of $R \leq 0$. In particular, M has infinite volume.

Lemma 5.2. *([Lin 2015b]) Let (M^n, g) be an n-dimensional complete Riemannian manifold. Then*

$$\mathrm{Ric} \geq -|T|g - \frac{|R|}{\sqrt{n}}g$$

in the sense of quadratic forms. Here T stands for the traceless tensor, namely

$$T = \mathrm{Ric} - \frac{R}{n}g.$$

Now, we introduce our result.

Theorem 5.3. *Let (M^n, g), $n \geq 3$, be an n-dimensional complete, simply connected, locally conformally flat Riemannian manifold. If one of the following conditions*

1. $$\|T\|_{n/2} + \frac{\|R\|_{n/2}}{\sqrt{n}} < \frac{4(Q-1+\kappa_p)}{SQ^2}.$$
2. *the scalar curvature R is nonpositive and*
$$K_{p,Q,n} := \frac{Q-1+\kappa_p}{Q^2} - \frac{n-1}{\sqrt{n}(n-2)} > 0$$
and
$$\|T\|_{n/2} < \frac{4K_{p,Q,n}}{S} = 4K_{p,Q,n}\mathcal{Y}(\mathbb{S}^n).$$

holds true, then every p-harmonic 1-form with finite $L^Q (Q \geq 2)$ norm on M is trivial. Here

$$\kappa_p = \min\left\{1, \frac{(p-1)^2}{n-1}\right\}.$$

Proof. Since M satisfies a Sobolev inequality, M must have infinite volume. Therefore, as in the previous part, we only need to prove that if ω is any p-harmonic 1-form with finite L^Q-norm then $\omega = 0$ in M_+. Now, applying the Bochner formula to the form $|\omega|^{p-2}\omega$, we have

$$\begin{aligned}\frac{1}{2}\Delta|\omega|^{2(p-1)} = &\left|\nabla\left(|\omega|^{p-2}\omega\right)\right|^2 - \left\langle \delta d(|\omega|^{p-2}\omega), |\omega|^{p-2}\omega\right\rangle \\ &+ |\omega|^{2(p-2)}\,\mathrm{Ric}(\omega,\omega).\end{aligned}$$

From this and Lemma 5.2 and the Kato type inequality (see, Lemma 2.2), we obtain

$$|\omega|\Delta|\omega|^{p-1} \geq \kappa_p|\omega|^{p-2}\left|\nabla|\omega|\right|^2 - \left\langle \delta d(|\omega|^{p-2}\omega), \omega\right\rangle - |T||\omega|^p - \frac{|R|}{\sqrt{n}}|\omega|^p.$$

We choose cutoff function $\varphi \in \mathcal{C}_0^\infty(M_+)$ as in the proof of Theorem 3.1. Multiplying both sides of the last inequality by $\varphi^2|\omega|^q$, $(q = Q-p)$ and then integrating by parts, we obtain

$$\begin{aligned}\int_{M_+} \varphi^2|\omega|^{q+1}\Delta|\omega|^{p-1} \geq &\kappa_p \int_{M_+} \varphi^2|\omega|^{p+q-2}\left|\nabla|\omega|\right|^2 - \int_{M_+}\left\langle \delta d(|\omega|^{p-2}\omega), \varphi^2|\omega|^q\omega\right\rangle \\ &- \int_{M_+}|T|\varphi^2|\omega|^{p+q} - \frac{1}{\sqrt{n}}\int_{M_+}|R|\varphi^2|\omega|^{p+q},\end{aligned}$$

and then,

$$\int_{M_+} \left\langle \nabla(\varphi^2|\omega|^{q+1}), \nabla|\omega|^{p-1} \right\rangle + \kappa_p \int_{M_+} \varphi^2 |\omega|^{p+q-2} \left|\nabla|\omega|\right|^2$$
$$\leq \int_{M_+} \left\langle d(|\omega|^{p-2}\omega), d(\varphi^2|\omega|^q\omega) \right\rangle + \int_{M_+} |T|\varphi^2|\omega|^{p+q} + \frac{1}{\sqrt{n}} \int_{M_+} |R|\varphi^2|\omega|^{p+q}.$$

By the hypotheses and Lemma 5.1, we obtain that, for some constant $S > 0$ and all functions $f \in C_0^\infty(M)$,

$$\left(\int_{M_+} f^{\frac{2n}{n-2}} dv \right)^{\frac{n-2}{n}} \leq S \int_{M_+} |\nabla f|^2 dv.$$

Using this and (3..10), we have that, for any $\varepsilon > 0$,

$$\begin{aligned}
\int_{M_+} |T|\varphi^2|\omega|^{p+q} \leq & \|T\|_{n/2} \left(\int_{M_+} \left(\varphi|\omega|^{(p+q)/2} \right)^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}} \\
\leq & S\|T\|_{n/2} \int_{M_+} \left| \nabla \left(\varphi|\omega|^{(p+q)/2} \right) \right|^2 \\
\leq & S\|T\|_{n/2}(1+\varepsilon)\frac{(p+q)^2}{4} \int_{M_+} \varphi^2|\omega|^{p+q-2} \left|\nabla|\omega|\right|^2 \\
& + S\|T\|_{n/2} \left(1 + \frac{1}{\varepsilon}\right) \int_{M_+} |\omega|^{p+q} |\nabla\varphi|^2. \qquad (5..2)
\end{aligned}$$

From (3..3), (3..4), (3..5) and (5..2) we obtain that, for any $\varepsilon > 0$,

$$\left[+q - 1 + \kappa_p - \varepsilon(2p-3) - S\|T\|_{n/2}(1+\varepsilon)\frac{(p+q)^2}{4} \right] \int_{M_+} \varphi^2|\omega|^{p+q-2} |\nabla|\omega||^2$$
$$\leq \left[\frac{\kappa_p}{\varepsilon} + S\|T\|_{n/2} \left(1 + \frac{1}{\varepsilon}\right) \right] \int_{M_+} |\omega|^{p+q} |\nabla\varphi|^2 + \frac{1}{\sqrt{n}} \int_{M_+} \varphi^2 |R||\omega|^{p+q}. \qquad (5..3)$$

For the last term in the right hand side of (5..3), we can estimate in the following two ways.

1. Similarly to (5..2), we have that, for any $\varepsilon > 0$,

$$\begin{aligned}
\int_{M_+} |R|\varphi^2|\omega|^{p+q} \leq & S\|R\|_{n/2}(1+\varepsilon)\frac{(p+q)^2}{4} \int_{M_+} \varphi^2|\omega|^{p+q-2} \left|\nabla|\omega|\right|^2 \\
& + S\|R\|_{n/2} \left(1 + \frac{1}{\varepsilon}\right) \int_{M_+} |\omega|^{p+q} |\nabla\varphi|^2.
\end{aligned}$$

Consequently, for any $\varepsilon > 0$,

$$C_\varepsilon \int_{M_+} \varphi^2|\omega|^{p+q-2} |\nabla|\omega||^2 \leq D_\varepsilon \int_{M_+} |\omega|^{p+q} |\nabla\varphi|^2,$$

where

$$\begin{aligned}
C_\varepsilon := & \, p + q - 1 + \kappa_p - \varepsilon(2p-3) - S(1+\varepsilon)\frac{(p+q)^2}{4} \left(\|T\|_{n/2} + \frac{\|R\|_{n/2}}{\sqrt{n}} \right) \\
= & \, Q - 1 + \kappa_p - \varepsilon(2p-3) - S(1+\varepsilon)\frac{Q^2}{4} \left(\|T\|_{n/2} + \frac{\|R\|_{n/2}}{\sqrt{n}} \right)
\end{aligned}$$

and

$$D_\varepsilon := \frac{2p-3}{\varepsilon} + S\left(1 + \frac{1}{\varepsilon}\right) \left(\|T\|_{n/2} + \frac{\|R\|_{n/2}}{\sqrt{n}} \right).$$

Since

$$\|T\|_{n/2}+\frac{\|R\|_{n/2}}{\sqrt{n}}<\frac{4(Q-1+(p-1)^2/(n-1))}{SQ^2},$$

there are some small enough $\varepsilon>0$ and constant $K=K(\varepsilon)>0$ such that

$$\frac{4}{Q^2}\int_{M_+}\left|\nabla|\omega|^{Q/2}\right|^2\varphi^2\leq K\int_{M_+}|\omega|^Q|\nabla\varphi|^2.$$

Using the same argument as in the proof of the first part of Theorem 3.1, this inequality implies that ω is zero.

2. If the scalar curvature R is nonpositive, then from (5..1) we have that

$$\int_{M_+}|R|f^2dv\leq\frac{4(n-1)}{n-2}\int_{M_+}|\nabla f|^2dv,\ \forall f\in C_0^\infty(M).$$

From this and (3..10), we have that, for any $\varepsilon>0$,

$$\begin{aligned}\int_{M_+}|R|\varphi^2|\omega|^{p+q}\leq&\frac{4(n-1)}{n-2}\int_{M_+}|\nabla(\varphi|\omega|^{(p+q)/2})|^2\\ \leq&\frac{4(n-1)}{n-2}(1+\varepsilon)\frac{(p+q)^2}{4}\int_{M_+}\varphi^2|\omega|^{p+q-2}\,|\nabla|\omega||^2\\ &+\frac{4(n-1)}{n-2}\left(1+\frac{1}{\varepsilon}\right)\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2.\end{aligned}$$

Substituting this inequality into (5..3) yields that, for any $\varepsilon>0$,

$$C_\varepsilon\int_{M_+}\varphi^2|\omega|^{p+q-2}|\nabla|\omega||^2\leq D_\varepsilon\int_{M_+}|\omega|^{p+q}|\nabla\varphi|^2,$$

where

$$\begin{aligned}C_\varepsilon:=&\,p+q-1+\kappa_p-\varepsilon(2p-3)-(1+\varepsilon)\frac{(p+q)^2}{4}\left(S\|T\|_{n/2}+\frac{4(n-1)}{(n-2)\sqrt{n}}\right)\\ =&\,Q-1+\kappa_p-\varepsilon(2p-3)-(1+\varepsilon)\frac{Q^2}{4}\left(S\|T\|_{n/2}+\frac{4(n-1)}{(n-2)\sqrt{n}}\right),\end{aligned}$$

and

$$D_\varepsilon:=\frac{2p-3}{\varepsilon}+\left(1+\frac{1}{\varepsilon}\right)\left(S\|T\|_{n/2}+\frac{4(n-1)}{(n-2)\sqrt{n}}\right).$$

Since

$$K_{p,Q,n}:=\frac{Q-1+\kappa_p}{Q^2}-\frac{n-1}{\sqrt{n}(n-2)}>0$$

and

$$\|T\|_{n/2}<\frac{4K_{p,Q,n}}{S},$$

there are some small enough $\varepsilon>0$ and constant $K=K(\varepsilon)>0$ such that

$$\frac{4}{Q^2}\int_{M_+}\left|\nabla|\omega|^{Q/2}\right|^2\varphi^2\leq K\int_{M_+}|\omega|^Q|\nabla\varphi|^2.$$

As in previous part, we infer that $\omega=0$. The proof is complete. □

Note that when $q=0$, we recover Theorem 1.3 and Theorem 1.4 in [Lin 2015b]. Hence, Theorem 5.3 can be considered as a generalization of Lin's results to the nonlinear setting. Now, we give a simple proof of Theorem 1.3.

Proof of Theorem 1.3. The proof follows by combining Theorem 3.2 and Theorem 5.3 with $Q=p$. □

Acknowledgment: A part of this paper was written during a stay of the authors at Vietnam Institute for Advanced Study in Mathematics (VIASM) and a stay of the first author at Institut Fourier, UMR 5582, Grenoble. We would to express our thanks to the staff there for the hospitality.

REFERENCES

[Calderbank et al. 2000] D. M. J. CALDERBANK, P. GAUDUCHON, AND M. HERZLICH, *Refined Kato inequalities and conformal weights in Riemannian geometry*, Jour. Funct. Anal., **173** (2000), 214 - 255. 2, 5

[Carron 1998] G. CARRON, *Une suite exacte en L^2-cohomologie*, Duke Math. Jour. **95** (1998), 343 - 372 1

[Carron 2007] G. CARRON, *L^2 harmonic forms on non-compact Riemannian manifolds*, Preprint 2007, see ArXiv:0704.3194v1. 1

[Chang et al. 2016] S. C. CHANG, J. T. CHEN, AND S. W. WEI, *Liouville properties for p-harmonic maps with finite q-energy*, Trans. Amer. Math. Soc., **368** (2016), 787 - 825. 2, 3, 12

[Chang et al. 2010] L. C. CHANG, C. L. GUO, AND C. J. A. SUNG, *p-harmonic 1-forms on complete manifolds*, Arch. Math., **94** (2010) 183 - 192. 2

[Chen and Sung 2009] J. T. R. CHEN, AND C. J. A. SUNG, *Harmonic forms on manifolds with weighted Poincaré inequality*, Pacific Jour. Math., **242** (2009) No. 2, 201 - 214. 3

[Cibotaru and Zhu 2012] D. CIBOTARU, AND P. ZHU, *Refined Kato inequalities for harmonic fields on Kähler manifolds*, Pacific Jour. Math., **256** (2012), No.1, 51 - 66. 2, 5

[Dung 2017] N. T. DUNG, *Vanishing theorems for p-harmonic ℓ-forms on Riemannian manifolds with a weighted Poincaré inequality*, Nonlinear Analysis: TMA., 150 (2017) 138 - 150. 3, 4

[Dung and Dat 2016] N. T. DUNG, AND N. D. DAT, *Local and global sharp gradient estimates for weighted p-harmonic functions*, Jour. Math. Anal. Appl., **443** (2016), 959 - 980. 4

[Dung and Seo 2017] N. T. DUNG, AND K. SEO, *p-harmonic functions and connectedness at infinity of complete Riemannian manifolds*, Annali di Matematica Pura ed Applicata, to appear. Doi:10.1007/s10231-016-0625-0. 2, 5

[Dung and Sung 2014] N. T. DUNG, AND C. J. A. SUNG, *Manifolds with a weighted Poincaré inequality*, Proc. Amer. Math. Soc., **142** (2014) No. 5, 1783 - 1794 3

[Duzaar and Fuchs 1990] F. DUZAAR, AND M. FUCHS, *On removable singularities of p-harmonic maps*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **7** (1990), 385 - 405. 9

[Hoffman and Spruck 1974] D. HOFFMAN, AND J. SPRUCK, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, Comm. Pure Appl. Math., **27** (1974), 715-727. 5

[Kotschwar and Ni 2009] B. KOTSCHWAR, AND L. NI, *Gradient estimate for p-harmonic function, $1/H$ flow and an entropy formula*, Ann. Sci. Éc. Norm. Supér., **42** (2009), 1-36 2

[Lam 2010] K. H. LAM, *Results on a weighted Poincaré inequality of complete manifolds.*, Trans. Amer. Math. Soc., **362** (2010), no. 10, 5043 - 5062 3

[Li 2012] P. LI, **Geometric analysis**, Cambridge Studies in Advanced Mathematics, 2012 5

[Li 1980] P. LI, *On the Sobolev constant and the p-spectrum of a compact Riemannian manifold*, Ann. Sci. École Norm. Sup. **13** (1980), 451-468. 1

[Li and Wang 2001] P. LI, AND J. WANG, *Complete manifolds with positive spectrum,* Jour. Diff. Geom., **58** (2001), 501 - 534. 3

[Li and Wang 2006] P. LI, AND J. WANG, *Weighted Poincaré inequality and rigidity of complete manifolds*, Ann. Sci. École Norm. Sup., **39** (2006), no. 6, 921 - 982. 3

[Lin 2015a] H. Z. LIN, *On the Structure of submanifolds in Euclidean space with flat normal bundle*, Results in Math., **68** (2015), Issue 3, pp 313-329. 1, 6

[Lin 2015b] H. Z. LIN, *On the structure of conformally flat Riemannian manifolds*, Nonlinear Analysis: TMA, **123-124** (2015), 115-125. 2, 17, 19

[Lin 2015c] H. Z. LIN, *L^2 harmonic forms on submanifolds in a Hadamard manifold*, Nonlinear Analysis: TMA, **125** (2015), 310-322. 2, 6

[Lin 2016] H. Z. LIN, *On the structure of submanifolds in the hyperbolic space*, Monatshefte für Mathematik, **180** (2016), Issue 3, 579 - 594 2

[Moser 2007] R. MOSER, *The inverse mean curvature flow and $p-$harmonic functions*, Jour. Eur. Math. Soc., **9** (2007), 77-83. 2

[Nakauchi 1998] N. NAKAUCHI, *A Liouville type theorem for p-harmonic maps*, Osaka J. Math. **35** (1998), no. 2, 303-312. 9

[Schoen and Yau 1994] R. SCHOEN, AND S.T. YAU, **Lectures on Differential Geometry**, Internatioal Press, 1994. 16

[Seo 2014] K. SEO, *L^p harmonic 1-forms and first eigenvalue of a stable minimal hypersurface*, Pacific Jour. Math., **268** (2014), no. 1, 205-229. 2

[Sung and Wang 2014] C. J. A. SUNG, AND J. WANG, *Sharp gradient estimate and spectral rigidity for p-Laplacian*, Math. Res. Lett., **21** (2014), no. 4, 885-904. 4

[Tanno 1996] S. TANNO, *L^2 harmonic forms and stability of minimal hypersurfaces*, Jour. Math. Soc. Japan, **48** (1996) 761- 768. 1

[Tolksdorff 1984] P. TOLKSDORF, *Regularity for a more general class of quasilinear elliptic equations*, Jour. Diff. Equa., **51** (1984), no. 1, 126-150. 12

[Vieira 2014] M. VIEIRA, *Vanishing theorems for L^2 harmonic forms on complete Riemannian manifolds*, Geom. Dedicata, **184** (2016), 175-191. 3, 14, 15

[Wan and Xin 2004] TOM Y. H. WAN, AND Y. L. XIN, *Vanishing Theorems for Conformally Compact Manifolds*, Comm. Partial Diff. Equa., **29** (2004), No. 7-8, 1267-1279. 2

[Wang 2002] X. D. WANG, *On the L^2-Cohomology of a convex cocompact hyperbolic manifold*, Duke Math. Jour., **115** (2002), No.2, 311-327. 2

[Zhang 2001] X. ZHANG, *A note on p-harmonic* 1*-forms on complete manifolds*, Canad. Math. Bull., **44** (2001) 376 - 384. 2

[Zhu 2011] P. ZHU, *Harmonic two forms on manifolds with nonnegative isotropic curvature*, Ann. Global Anal. Geom., **40** (2011), 427 - 434. 1, 2

[Zhu 2017] P. ZHU, *Rigidity of Complete Minimal Hypersurfaces in the Euclidean Space*, Results. Math., **71** (2017), 63 - 71. 1

[Zhu 2001] P. ZHU, *L^2-harmonic forms and stable hypersurfaces in space forms*, Arch. der Math., **97** (2001), 271-279.

(N.T. Dung) DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, VIÊT NAM NATIONAL UNIVERSITY, HA NÔI, VIÊT NAM

E-mail address: dungmath@gmail.com

(P.T. Tien) DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, VIÊT NAM NATIONAL UNIVERSITY, HA NÔI, VIÊT NAM

E-mail address: phamtien@mail.ru, phamtien@vnu.edu.vn